

An Actinide Dependent Shadow Biosphere of Archaea and Viroids in Neuropsychiatric Disorders

Ravikumar Kurup A.^{1,*}; Parameswara Achutha Kurup¹

¹Professor of Metabolic Medicine and Neurology

The Metabolic Disorders Research Centre, TC 4/1525, Gouri Sadan, Kattu Road, North of Cliff House, Kowdiar PO, Trivandrum, Kerala, India

*Corresponding author.

Email: ravikurup13@yahoo.in

Received 17 September 2011; accepted 19 November 2011

Abstract

Objective: Endogenous digoxin has been related to the pathogenesis of schizophrenia, autism and primary seizure disorder. The possibility of endogenous digoxin synthesis by actinide based primitive organism like archaea with a mevalonate pathway and cholesterol catabolism was considered.

Methods: 10 cases each of schizophrenia, autism and primary seizure disorder before starting treatment and 10 age and sex matched healthy controls from general population were chosen for the study. Cholesterol substrate was added to the plasma of the patients and the generation of cytochrome F420, free RNA, free DNA, polycyclic aromatic hydrocarbon, hydrogen peroxide, serotonin, pyruvate, ammonia, glutamate, cytochrome C, hexokinase, ATP synthase, HMG CoA reductase, digoxin and bile acids were studied. The changes with the addition of antibiotics and rutilite to the patient's plasma were also studied. The statistical analysis was done by ANOVA.

Results: The parameters mentioned above were increased the patient's plasma with addition of cholesterol substrate. The addition of antibiotics to the patient's plasma caused a decrease in all the parameters while addition of rutilite increased their levels.

Conclusion: An actinide dependent shadow biosphere of archaea and viroids is described in schizophrenia, autism and primary seizure disorder contributing to their pathogenesis.

Key words: Schizophrenia; Autism; Primary seizure disorder; Archaea; Viroids; Cholesterol; Actinide

Ravikumar Kurup A., Parameswara Achutha Kurup (2011). An Actinide Dependent Shadow Biosphere of Archaea and Viroids in Neuropsychiatric Disorders. *Advances in Natural Science*, 4(2), 52-58. Available from: URL: <http://www.cscanada.net/index.php/css/article/view/j.ans.1715787020110402.267> DOI: <http://dx.doi.org/10.3968/j.ans.1715787020110402.267>.

INTRODUCTION

Endomyocardial fibrosis (EMF) along with the root wilt disease of coconut is endemic to Kerala with its radioactive actinide beach sands. Actinides like rutile producing intracellular magnesium deficiency due to rutile-magnesium exchange sites in the cell membrane has been implicated in the etiology of EMF^[1]. Endogenous digoxin, a steroidal glycoside which functions as a membrane sodium-potassium ATPase inhibitor has also been related to its etiology due to the intracellular magnesium deficiency it produces^[2]. Organisms like phytoplasmata and viroids have also been demonstrated to play a role in the etiology of these diseases^[3,4]. Endogenous digoxin has been related to the pathogenesis of schizophrenia, autism and primary seizure disorder^[2]. The possibility of endogenous digoxin synthesis by actinide based primitive organism like archaea with a mevalonate pathway and cholesterol catabolism was considered^[5,6,7]. Davies has put forward the concept of a shadow biosphere of organisms with alternate biochemistry present in earth itself^[8]. An actinide dependent shadow biosphere of archaea and viroids in the above mentioned disease states is described^[6].

MATERIALS AND METHODS

Informed consent of the subjects and the approval of the ethics committee were obtained for the study. The following groups were included in the study: schizophrenia, autism and primary seizure disorder/

primary generalized epilepsy. There were 10 patients in each group and each patient had an age and sex matched healthy control selected randomly from the general population. The blood samples were drawn in the fasting state before treatment was initiated. Plasma from fasting heparinised blood was used and the experimental protocol was as follows (I) Plasma+phosphate buffered saline, (II) same as I+cholesterol substrate, (III) same as II+rutile 0.1 mg/ml, (IV) same as II+ciprofloxacin and doxycycline each in a concentration of 1 mg/ml. Cholesterol substrate was prepared as described by Richmond^[9]. Aliquots were withdrawn at zero time immediately after mixing and after incubation at 37°C for 1 hour. The following estimations were carried out:- Cytochrome F420, free RNA, free DNA, muramic acid, polycyclic aromatic hydrocarbon, hydrogen peroxide, dopamine, pyruvate, ammonia, glutamate, cytochrome C, hexokinase, ATP synthase, HMG CoA reductase, digoxin and bile acids^[10,11,12,13]. Cytochrome F420 was estimated fluorimetrically (excitation wavelength 420 nm and emission wavelength 520 nm). Polycyclic aromatic hydrocarbon was estimated by measuring hydrogen peroxide liberated by using glucose reagent. The statistical analysis was done by

ANOVA.

RESULTS

The parameters checked as indicated above were:- cytochrome F420, free RNA, free DNA, muramic acid, polycyclic aromatic hydrocarbon, hydrogen peroxide, serotonin, pyruvate, ammonia, glutamate, cytochrome C, hexokinase, ATP synthase, HMG CoA reductase, digoxin and bile acids. Plasma of control subjects showed increased levels of the above mentioned parameters with after incubation for 1 hour and addition of cholesterol substrate resulted in still further significant increase in these parameters. The plasma of patients showed similar results but the extent of increase was more. The addition of antibiotics to the control plasma caused a decrease in all the parameters while addition of rutile increased their levels. The addition of antibiotics to the patient's plasma caused a decrease in all the parameters while addition of rutile increased their levels but the extent of change was more in patient's sera as compared to controls. The results are expressed in tables 1-7 as percentage change in the parameters after 1 hour incubation as compared to the values at zero time.

Table 1
Effect of Rutile and Antibiotics on Muramic Acid and Dopamine

Group	DOPAMINE % (Increase with Rutile)		DOPAMINE % (Decrease with Doxy)		Muramic acid % change (Increase with Rutile)		Muramic acid % change (Decrease with Doxy+Cipro)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.41	0.15	18.63	0.12	4.34	0.15	18.24	0.37
Schizo	21.88	1.19	66.28	3.60	23.02	1.65	67.61	2.77
Seizure	22.29	1.33	65.38	3.62	22.13	2.14	66.26	3.93
Autism	22.76	2.20	67.63	3.52	22.79	2.20	64.26	6.02
F value	403.394		680.284		348.867		364.999	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

Table 2
Effect of Rutile and Antibiotics on Free DNA and RNA

Group	DNA % change (Increase with Rutile)		DNA % change (Decrease with Doxy)		RNA % change (Increase with Rutile)		RNA % change (Decrease with Doxy)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.37	0.15	18.39	0.38	4.37	0.13	18.38	0.48
Schizo	23.28	1.70	61.41	3.36	23.59	1.83	65.69	3.94
Seizure	23.40	1.51	63.68	4.66	23.08	1.87	65.09	3.48
Autism	22.12	2.44	63.69	5.14	23.33	1.35	66.83	3.27
F value	337.577		356.621		427.828		654.453	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

Table 3
Effect of Rutile and Antibiotics on HMG CoA Reductase and PAH

Group	HMG CoA R % change (Increase with Rutile)		HMG CoA R % change (Decrease with Doxy)		PAH % change (Increase with Rutile)		PAH % change (Decrease with Doxy)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.30	0.20	18.35	0.35	4.45	0.14	18.25	0.72
Schizo	22.91	1.92	61.63	6.79	23.01	1.69	59.49	4.30
Seizure	23.09	1.69	61.62	8.69	22.67	2.29	57.69	5.29
Autism	22.72	1.89	64.51	5.73	22.61	1.42	64.48	6.90
F value	319.332		199.553		391.318		257.996	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

Table 4
Effect of Rutile and Antibiotics on Digoxin and Bile Acids

Group	Digoxin (ng/ml) (Increase with Rutile)		Digoxin (ng/ml) (Decrease with Doxy+Cipro)		Bile acids % change (Increase with Rutile)		Bile acids % change (Decrease with Doxy)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	0.11	0.00	0.054	0.003	4.29	0.18	18.15	0.58
Schizo	0.55	0.06	0.219	0.043	23.20	1.87	57.04	4.27
Seizure	0.51	0.05	0.199	0.027	22.61	2.22	66.62	4.99
Autism	0.53	0.08	0.205	0.041	22.21	2.04	63.84	6.16
F value	135.116		71.706		290.441		203.651	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

Table 5
Effect of Rutile and Antibiotics on Pyruvate and Hexokinase

Group	Pyruvate % change (Increase with Rutile)		Pyruvate % change (Decrease with Doxy)		Hexokinase % change (Increase with Rutile)		Hexokinase % change (Decrease with Doxy)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.34	0.21	18.43	0.82	4.21	0.16	18.56	0.76
Schizo	20.99	1.46	61.23	9.73	23.01	2.61	65.87	5.27
Seizure	20.94	1.54	62.76	8.52	23.33	1.79	62.50	5.56
Autism	21.91	1.71	58.45	6.66	22.88	1.87	65.45	5.08
F value	321.255		115.242		292.065		317.966	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

Table 6
Effect of Rutile and Antibiotics on Hydrogen Peroxide and Delta Amino Levulinic Acid

Group	H ₂ O ₂ % (Increase with Rutile)		H ₂ O ₂ % (Decrease with Doxy)		ALA % (Increase with Rutile)		ALA % (Decrease with Doxy)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.43	0.19	18.13	0.63	4.40	0.10	18.48	0.39
Schizo	22.50	1.66	60.21	7.42	22.52	1.90	66.39	4.20
Seizure	23.81	1.19	61.08	7.38	22.83	1.90	67.23	3.45
Autism	23.52	1.49	63.24	7.36	23.20	1.57	66.65	4.26
F value	380.721		171.228		372.716		556.411	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

Table 7
Effect of Rutile and Antibiotics on ATP Synthase and Cytochrome F 420

Group	ATP synthase % (Increase with Rutile)		ATP synthase % (Decrease with Doxy)		CYT F420 % (Increase with Rutile)		CYT F420 % (Decrease with Doxy)	
	Mean	± SD	Mean	± SD	Mean	± SD	Mean	± SD
Normal	4.40	0.11	18.78	0.11	4.48	0.15	18.24	0.66
Schizo	23.67	1.42	67.39	3.13	23.24	2.01	58.72	7.08
Seizure	23.09	1.90	66.15	4.09	23.46	1.87	59.27	8.86
Autism	22.60	1.64	66.86	4.21	21.68	1.90	57.93	9.64
F value	449.503		673.081		306.749		130.054	
P value	< 0.001		< 0.001		< 0.001		< 0.001	

DISCUSSION

There was increase in cytochrome F420 indicating archaeal growth. The archaea can synthesise and use cholesterol as a carbon and energy source^[14,15]. The archaeal origin of the enzyme activities was indicated by antibiotic induced suppression. The study indicates the presence of actinide based archaea with an alternate actinide based enzymes or metalloenzymes in the system as indicated by rutile induced increase in enzyme activities^[16]. There was also an increase in archaeal HMG CoA reductase activity indicating increased cholesterol synthesis by the archaeal mevalonate pathway. The archaeal beta hydroxyl steroid dehydrogenase activity indicating digoxin synthesis and archaeal cholesterol hydroxylase activity indicating bile acid synthesis were increased^[17]. The archaeal cholesterol oxidase activity was increased resulting in generation of pyruvate and hydrogen peroxide^[15]. The pyruvate gets converted to glutamate and ammonia by the GABA shunt pathway. The archaeal aromatization of cholesterol generating PAH, serotonin and dopamine was also detected^[17]. The archaeal glycolytic hexokinase activity and archaeal extracellular ATP synthase activity were increased. The archaea can undergo magnetite and calcium carbonate mineralization and can exist as calcified nanoforms^[18]. There was an increase in free RNA indicating self replicating RNA viroids and free DNA indicating generation of viroid complementary DNA strands by archaeal reverse transcriptase activity. The actinides modulate RNA folding and catalyse its ribozymal action. Digoxin can cut and paste the viroidal strands by modulating RNA splicing generating RNA viroidal diversity. The viroids are evolutionarily escaped archaeal group I introns which have retrotransposition and self splicing qualities^[19]. Archaeal pyruvate can produce histone deacetylase inhibition resulting in endogenous retroviral (HERV) reverse transcriptase and integrase expression. This can integrate the RNA viroidal complementary DNA into the noncoding region of eukaryotic non coding DNA using HERV integrase as has been described for borna and ebola viruses^[20]. The noncoding DNA is lengthened by integrating RNA viroidal complementary DNA with the integration going

on as a continuing event. The archaea genome can also get integrated into human genome using integrase as has been described for trypanosomes^[21]. The integrated viroids and archaea can undergo vertical transmission and can exist as genomic parasites^[20,21]. This increases the length and alters the grammar of the noncoding region producing memes or memory of acquired characters^[22]. The viroidal complementary DNA can function as jumping genes producing a dynamic genome important in storage of synaptic information, HLA gene expression and neurodevelopmental gene expression. The alteration in DNA sequences produced by viroidal complementary DNA jumping genes can lead onto schizophrenia, autism and primary seizure disorder. The RNA viroids can regulate mrna function by RNA interference^[19]. The phenomena of RNA interference can modulate T cell and B cell function, neuronal transmission and euchromatin/ heterochromatin expression. The RNA viroid induced mRNA interference can modulate dopaminergic, glutamatergic and serotonergic synaptic transmission contributing to the pathogenesis of schizophrenia, autism and primary seizure disorder.

The presence of muramic acid, HMG CoA reductase and cholesterol oxidase activity inhibited by antibiotics indicates the presence of bacteria with mevalonate pathway. The bacterial with mevalonate pathway include streptococcus, staphylococcus, actinomycetes, listeria, coxiella and borrelia^[23]. The bacteria and archaea with mevalonate pathway and cholesterol catabolism had a evolutionarily advantage and constitutes the isoprenoidal clade organism with the archaea evolving into mevalonate pathway gram positive and gram negative organism through horizontal gene transfer of viroidal and virus genes^[24]. The isoprenoidal clade prokaryotes develop into other groups of prokaryotes via viroidal/virus as well as eukaryotic horizontal gene transfer producing bacterial speciation^[25]. The RNA viroids and its complementary DNA developed into cholesterol enveloped RNA and DNA viruses like herpes, retrovirus, influenza virus, borna virus, cytomegalo virus and Ebstein Barr virus by recombining with eukaryotic and human genes resulting in viral speciation. Bacterial and viral species are ill defined and fuzzy with all of them forming one common

genetic pool with frequent horizontal gene transfer and recombination. Thus the multi and unicellular eukaryote with its genes serves the purpose of prokaryotic and viral speciation. The multicellular eukaryote developed so that their endosymbiotic archaeal colonies could survive and forage better. The multicellular eukaryotes are like bacterial biofilms. The archaea and bacteria with a mevalonate pathway uses the extracellular RNA viroids and DNA viroids for quorum sensing and in the generation of symbiotic biofilm like structures which develop into multicellular eukaryotes^[26,27]. The endosymbiotic archaea and bacteria with mevalonate pathway still uses the RNA viroids and DNA viroids for the regulation of multicellular eukaryote. Pollution is induced by the primitive nanoarchaea and mevalonate pathway bacteria synthesised PAH and methane leading on to redox stress. Redox stress leads to sodium potassium ATPase inhibition, inward movement of plasma membrane cholesterol, defective SREBP sensing, increased cholesterol synthesis and nanoarchaeal/mevalonate pathway bacterial growth^[28]. Redox stress leads on to viroidal and archaeal multiplication. Redox stress can also lead to HERV reverse transcriptase and integrase expression. The noncoding DNA is formed of integrating RNA viroidal complementary DNA and archaea with the integration going on as a continuing event. The archaeal pox like dsDNA virus forms evolutionarily the nucleus. The integrated viroidal, archaeal and mevalonate pathway bacterial sequences can undergo vertical transmission and can exist as genomic parasites. The genomic integrated archaea, mevalonate pathway bacteria and viroids form a genomic reserve of bacteria and viruses which can recombine with human and eukaryotic genes producing bacterial and viral speciation. Bacteria and viruses have been related to the pathogenesis of schizophrenia, autism and primary seizure disorder. *Borrelia*, *Toxoplasma*, *Chlamydia*, *Mycoplasma*, retroviruses, herpes virus, influenza virus and borna virus contribute to the neuropathogenesis of schizophrenia. Clostridia, rubella virus, influenza virus and *Borrelia* are implicated in autism^[29,30,31]. Herpes virus is implicated in the pathogenesis of primary seizure disorder. The change in the length and grammar of the noncoding region produces eukaryotic speciation and individuality³². Changes in the length of noncoding region can lead onto disorders of consciousness like schizophrenia³³. The human endogenous retroviruses and change in the length and grammar of the noncoding region has been described in schizophrenia. The integration of nanoarchaea, mevalonate pathway prokaryotes and viroids in to the eukaryotic and human genome produces a chimera which can multiply producing biofilm like multicellular structures having a mixed archaeal, viroidal, prokaryotic and eukaryotic characters which is a regression from the multicellular eukaryotic tissue. This results in a new neuronal, metabolic, immune and tissue phenotype leading

to human diseases like schizophrenia, autism and primary seizure disorder. The microchimeras formed can lead to polyploidy. Neuronal polyploidy and microchimeras have been described in schizophrenia and autism.

The archaea and viroids can regulate the nervous system including the NMDA/GABA thalamocorticothalamic pathway mediating conscious perception^[2,34]. NMDA/GABA receptors can be modulated by digoxin induced calcium oscillations resulting in NMDA/glutamic acid decarboxylase (GAD) activity induction, PAH increasing NMDA activity and inducing GAD as well as viroid induced RNA interference modulating NMDA/GABA receptors^[2]. The cholesterol ring oxidase generated pyruvate can be converted by the GABA shunt pathway to glutamate and GABA. Increased NMDA transmission has been described in schizophrenia, autism and primary seizure disorder. The dipolar PAH and archaeal magnetite in the setting of digoxin induced sodium potassium ATPase inhibition can produce a pumped phonon system mediated frohlich model superconducting state inducing quantal perception with nanoarchaeal sensed gravity producing the orchestrated reduction of the quantal possibilities to the macroscopic world^[2,34]. Schizophrenia and autism are described as a disorder of consciousness and increased integration of archaea and viroids into the genome can contribute to its neuropathogenesis. The archaea can regulate limbic lobe transmission with archaeal cholesterol aromatase/ring oxidase generated norepinephrine, dopamine, serotonin and acetyl choline^[17]. Increased dopaminergic and serotonergic transmission is important in the pathogenesis of schizophrenia and autism. The higher degree of integration of the archaea into the genome produces increased digoxin synthesis producing right hemispheric dominance and lesser degree producing left hemispheric dominance^[2]. Right hemispheric dominance has been described in schizophrenia, autism and primary seizure disorder. The increased integration of archaea into the neuronal genome can produce increased cholesterol oxidase and aromatase mediated monoamine and NMDA transmission producing schizophrenia, autism and primary seizure disorder. The archaeal bile acids are chemically diverse and structurally different from human bile acids. The archaeal bile acids can bind olfactory GPCR receptors and stimulate the limbic lobe producing a sense of social identity. The dominance of archaeal bile acids over human bile acids in stimulating the olfactory GPCR-limbic lobe pathway leads to loss of social identity and schizophrenia/ autism^[35]. Archaea and RNA viroid can bind the TLR receptor induce NF κ B producing immune activation and cytokine TNF alpha secretion. The archaeal DXP and mevalonate pathway metabolites can bind $\gamma\delta$ TCR and digoxin induced calcium signaling can activate NF κ B producing chronic immune activation^[2,36]. The archaea and viroid induced chronic immune activation and generation of superantigens can lead on to autoimmune

disease. Immune activation and autoimmunity is important in the pathogenesis of schizophrenia, autism and primary seizure disorder. Autoantibodies have been described in schizophrenia, autism and primary seizure disorder. Archaea, viroids and digoxin can induce the host AKT PI3K, AMPK, HIF alpha and NFkB producing the Warburg metabolic phenotype^[37]. The increased glycolytic hexokinase activity, decrease in blood ATP, leakage of cytochrome C, increase in serum pyruvate and decrease in acetyl CoA indicates the generation of the Warburg phenotype. There is induction of glycolysis, inhibition of PDH activity and mitochondrial dysfunction resulting in inefficient energetics. Mitochondrial dysfunction can lead onto NMDA excitotoxicity and cell death important in schizophrenia and primary seizure disorder. Cholesterol oxidase activity, increased glycolysis related NADPH oxidase activity and mitochondrial dysfunction generates free radicals important in the pathogenesis of schizophrenia, autism and primary seizure disorder. The accumulated pyruvate enters the GABA shunt pathway and is converted to citrate which is acted upon by citrate lyase and converted to acetyl CoA, used for cholesterol synthesis^[37]. The pyruvate can be converted to glutamate and ammonia which is oxidised by archaea for energy needs. The increased cholesterol substrate leads to increased archaeal growth and digoxin synthesis leading to metabolic channeling to the mevalonate pathway. Hyperdigoxinemia is important in the pathogenesis of schizophrenia, autism and primary seizure disorder^[2]. The Warburg phenotype can contribute to the pathogenesis of schizophrenia by augmenting the bacterial shikimic acid pathway. The upregulated glycolysis consequent to the Warburg phenotype produces phosphoenolpyruvate, a basic substrate for the bacterial shikimic acid pathway which can synthesise monoamines and neuroactive alkaloids. The shikimic acid pathway can generate dopamine and serotonin producing the increased monoaminergic transmission in schizophrenia. The shikimic acid pathway can also synthesise the neuroactive alkaloids strychnine, nicotine, morphine, mescaline and LSD important in the pathogenesis of schizophrenia and autism. Endogenous neuroactive alkaloids has been described in schizophrenia, autism and primary seizure disorder by several workers^[2]. The upregulated glycolysis can also contribute to increased NMDA and GABA transmission in the thalamocorticothalamic pathway. The glycolytic pathway produces phosphoglycerate which is converted to phosphoserine and then serine which activates the NMDA receptor. The glycolytic enzyme glyceraldehyde-3-phosphate dehydrogenase is a GABA receptor kinase and activates GABA transmission. Thus the archaea and viroid induced Warburg phenotype can contribute to the pathogenesis of schizophrenia, autism and primary seizure disorder. The archaeal cholesterol catabolism can deplete the cell membranes of cholesterol resulting in alteration in lipid microdomains and their

related neurotransmitter receptor contributing to the altered NMDA, serotonergic and dopaminergic transmission in schizophrenia, autism and primary seizure disorder.

REFERENCES

- [1] Valiathan, M. S., Somers, K., & Kartha, C. C. (1993). *Endomyocardial Fibrosis*. Delhi: Oxford University Press.
- [2] Kurup, R., & Kurup, P. A. (2009). *Hypothalamic Digoxin, Cerebral Dominance and Brain Function in Health and Diseases*. New York: Nova Science Publishers.
- [3] Hanold, D., & Randies, J. W. (1991). Coconut Cadang-Cadang Disease and its Viroid Agent. *Plant Disease*, 75, 330-335.
- [4] Edwin, B. T., & Mohankumaran, C. (2007). Kerala wilt Disease Phytoplasma: Phylogenetic Analysis and Identification of a Vector, Proutista Moesta. *Physiological and Molecular Plant Pathology*, 71(1-3), 41-47.
- [5] Eckburg, P. B., Lepp, P.W., & Relman, D. A. (2003). Archaea and their Potential Role in Human Disease. *Infect Immun*, 71, 591-596.
- [6] Adam, Z. (2007). Actinides and Life's Origins. *Astrobiology*, 7, 6-10.
- [7] Schoner, W. (2002). Endogenous Cardiac Glycosides, A New Class of Steroid Hormones. *Eur J Biochem*, 269, 2440-2448.
- [8] Davies, P. C. W., Benner, S. A., Cleland, C. E., Lineweaver, C. H., McKay, C. P., & Wolfe-Simon, F. (2009). Signatures of a Shadow Biosphere. *Astrobiology*, 10, 241-249.
- [9] Richmond, W. (1973). Preparation and Properties of a Cholesterol Oxidase from Nocardia Species and its Application to the Enzymatic Assay of Total Cholesterol in Serum. *Clin Chem*, 19, 1350-1356.
- [10] Snell, E. D., & Snell, C. T. (1961). *Colorimetric Methods of Analysis* (Vol 3A). New York: Van Nostrand.
- [11] Glick, D. (1971). *Methods of Biochemical Analysis* (Vol 5). New York: Interscience Publishers.
- [12] Colowick, Kaplan, N.O. (1955). *Methods in Enzymology* (Vol 2). New York: Academic Press.
- [13] Maarten, A. H., Marie-Jose, M., Cornelia, G., van Helden-Meewsen, Fritz, E., & Marten, P. H. (1995). Detection of Muramic Acid in Human Spleen. *Infection and Immunity*, 63(5), 1652 – 1657.
- [14] Smit, A., & Mushegian, A. (2000). Biosynthesis of Isoprenoids via Mevalonate in Archaea: the Lost Pathway. *Genome Res*, 10(10), 1468-84.
- [15] Van der Geize, R., Yam, K., Heuser, T., Wilbrink, M. H., Hara, H., & Anderton, M. C. (2007). A Gene Cluster Encoding Cholesterol Catabolism in a Soil Actinomycete Provides Insight into Mycobacterium Tuberculosis Survival In Macrophages. *Proc Natl Acad Sci USA*, 104(6), 1947-52.
- [16] Francis, A. J. (1998). Biotransformation of Uranium and Other Actinides In Radioactive Wastes. *Journal of Alloys and Compounds*, 271(273), 78-84.
- [17] Probian, C., Wülfing, A., & Harder, J. (2003).

- Anaerobic Mineralization of Quaternary Carbon Atoms: Isolation of Denitrifying Bacteria on Pivalic Acid (2,2-Dimethylpropionic acid). *Applied and Environmental Microbiology*, 69(3), 1866-1870.
- [18] Vainshtein, M., Suzina, N., Kudryashova, E., & Ariskina, E. (2002). New Magnet-Sensitive Structures in Bacterial and Archaeal Cells. *Biol Cell*, 94(1), 29-35.
- [19] Tsagris, E. M., de Alba, A. E., Gozmanova, M., & Kalantidis, K. (2008). *Viroids, Cell Microbiol*, 10, 2168.
- [20] Horie, M., Honda, T., Suzuki, Y., Kobayashi, Y., Daito, T., & Oshida, T. (2010). Endogenous Non-Retroviral RNA Virus Elements in Mammalian Genomes. *Nature*, 463, 84-87.
- [21] Hecht, M., Nitz, N., Araujo, P., Sousa, A., Rosa, A., & Gomes, D. (2010). Genes from Chagas Parasite can Transfer to Humans and be Passed on to Children. Inheritance of DNA Transferred from American Trypanosomes to Human Hosts. *PLoS ONE*, 5, 2-10.
- [22] Flam, F. (1994). Hints of a Language in Junk DNA. *Science*, 266, 1320.
- [23] Horbach, S., Sahn, H., & Welle, R. (1993). Isoprenoid Biosynthesis in Bacteria: Two Different Pathways? *FEMS Microbiol Lett*, 111, 135-140.
- [24] Gupta, R. S. (1998). Protein Phylogenetics and Signature Sequences: A Reappraisal of Evolutionary Relationship Among Archaeobacteria, Eubacteria, and Eukaryotes. *Microbiol Mol Biol Rev*, 62, 1435-1491.
- [25] Hanage W., Fraser, C., & Spratt, B. (2005). Fuzzy Species Among Recombinogenic Bacteria. *BMC Biology*, 3, 6-10.
- [26] Webb J. S., Givskov, M., & Kjelleberg, S. (2003). Bacterial Biofilms: Prokaryotic Adventures in Multicellularity. *Curr Opin Microbiol*, 6(6), 578-85.
- [27] Whitchurch, C. B., Tolker-Nielsen, T., Ragas, P. C., & Mattick, J. S. (2002). Extracellular DNA Required for Bacterial Biofilm Formation. *Science*, 295(5559), 1487.
- [28] Chen, Y., Cai, T., Wang, H., Li, Z., Loreaux, E., & Lingrel, J. B. (2009). Regulation of Intracellular Cholesterol Distribution by Na/K-ATPase. *J Biol Chem*, 284(22), 14881-90.
- [29] Fritzsche, M. (2002). Seasonal correlation of sporadic schizophrenia to Ixodes ticks and Lyme borreliosis. *Int J Health Geogr*, 1(1), 2.
- [30] Waltrip, R. W. 2nd, Buchanan, R. W., Summerfelt, A., Breier, A., Carpenter, W. T. Jr., Bryant, N. L., Rubin, S. A., & Carbone, K. M. (1995). Borna Disease Virus and Schizophrenia. *Psych Res*, 56(1), 33-44.
- [31] Torrey, E. F., & Yolken, R. H. (2003). Toxoplasma Gondii and Schizophrenia. *Emerg Infect Dis*, 9(11), 1375-80.
- [32] Poole, A. M. (2006). Did group II Intron Proliferation in an Endosymbiont-Bearing Archaeon Create Eukaryotes? *Biol Direct*, 1, 36-40.
- [33] Villarreal, L. P. (2006). How Viruses Shape the Tree of Life. *Future Virology*, 1(5), 587-595.
- [34] Lockwood, M. (1989). *Mind, Brain and the Quantum*. Oxford: B. Blackwell.
- [35] Lefebvre, P., Cariou, B., Lien, F., Kuipers, F., & Staels, B. (2009). Role of Bile Acids and Bile Acid Receptors in Metabolic Regulation. *Physiol Rev*, 89(1), 147 - 191.
- [36] Eberl, M., Hintz, M., Reichenberg, A., Kollas, A., Wiesner, J., & Jomaa, H. (2010). Microbial Isoprenoid Biosynthesis and Human $\gamma\delta$ T Cell Activation. *FEBS Letters*, 544(1), 4-10.
- [37] Wallace, D. C. (2005). Mitochondria and Cancer: Warburg Addressed. *Cold Spring Harbor Symposia on Quantitative Biology*, 70, 363-374.