

A Fast Convergent Error Bound for Gaussian Interpolation

Lin-Tian Luh^{1,*}

Abstract: It's well known that there is a so-called exponential-type error bound for Gaussian interpolation which is the most powerful error bound hitherto. It's of the form $|f(x) - s(x)| \leq c_1(c_2d)^{\frac{c_3}{d}} \|f\|_h$ where f and s are the interpolated and interpolating functions respectively, c_1, c_2, c_3 are positive constants, d is the fill distance which roughly speaking measures the spacing of the data points, and $\|f\|_h$ is the h -norm of f where h is the Gaussian function. The error bound is suitable for $x \in R^n$, $n \geq 1$, and gets small rapidly as $d \rightarrow 0$. The drawback is that the crucial constants c_2 and c_3 get worse rapidly as n increases in the sense $c_2 \rightarrow \infty$ and $c_3 \rightarrow 0$ as $n \rightarrow \infty$. In this paper we raise an error bound of the form

$$|f(x) - s(x)| \leq c'_1(c'_2d)^{\frac{c'_3}{d}} \sqrt{d} \|f\|_h,$$

where c'_2 and c'_3 are independent of the dimension n . Moreover, $c'_2 \ll c_2$, $c_3 \ll c'_3$, and c'_1 is only slightly different from c_1 . What's important is that all constants c'_1, c'_2 and c'_3 can be computed without slight difficulty.

Key Words: Radial basis function; Interpolation; Error bound; Gaussian

1. INTRODUCTION

First, let h be a continuous radial function on R^n which is conditionally positive definite of order m . Given data points (x_j, f_j) , $j = 1, \dots, N$, where $X = \{x_1, \dots, x_N\}$ is a subset of points in R^n and the f_j are real or complex numbers, the so-called h spline interpolant of these data points is the function s defined by

$$s(x) = p(x) + \sum_{j=1}^N c_j h(x - x_j), \quad (1)$$

where $p(x)$ is a polynomial in P_{m-1}^n and c_j are chosen so that

$$\sum_{j=1}^N c_j q(x_j) = 0 \quad (2)$$

for all polynomials q in P_{m-1}^n and

$$p(x_i) + \sum_{j=1}^N c_j h(x_i - x_j) = f_i, \quad i = 1, \dots, N. \quad (3)$$

¹Department of Mathematics, Providence University Shalu, Taichung, Taiwan.
Email addresses: ltluh@pu.edu.tw.

*Corresponding author.

[†]Received 3 December 2010; accepted 15 February 2011.

Here P_{m-1}^n denotes the class of those polynomials of R^n of degree $\leq m-1$.

A famous property is that the system of equations (2) and (3) has a unique solution whenever X is a determining set for P_{m-1}^n and h is strictly conditionally positive definite. More details can be seen in [8]. Therefore in our case the interpolant $s(x)$ is well defined.

We clarify that X is said to be a determining set for P_{m-1}^n if X does not lie on the zero set of any nontrivial polynomial in P_{m-1}^n .

In this paper the function h is defined by

$$h(x) := e^{-\beta|x|^2}, \beta > 0. \quad (4)$$

This is the so-called Gaussian function.

In [10] Madych and Nelson raise the famous exponential-type error bound for the scattered data interpolation of Gaussian function, as mentioned in the abstract. The computation of the constants c_1, c_2 and c_3 can be found in [?]. This error bound is very powerful. However, too many data points may lead to a large condition number when solving the linear system (2) and (3). There will be a significant improvement of the ill-conditioning if a better error bound can be obtained so that a satisfactory error estimate is reached before too many data points are involved. This is what we are pursuing in this paper.

1.1 Polynomials and Simplices

In this paper we will use P_l^n to denote the space of polynomials of degree l in n variables. It's well known that it has dimension $\dim P_l^n = \binom{n+l}{n}$. We will denote $\dim P_l^n$ by N in this section. Let $E \subseteq R^n$ be compact.

The interpolation theory tells us that if $x_1, \dots, x_N \in E$ and do not lie on the zero set of any nontrivial $q \in P_l^n$, there exists Lagrange polynomials $l_i, i = 1, \dots, N$, of degree l defined by $l_i(x_j) = \delta_{ij}, 1 \leq i, j \leq N$, such that for any $f \in C(E)$, $(\Pi_l f)(x) := \sum_{i=1}^N f(x_i)l_i(x)$ is its interpolating polynomial. It's easily seen that $\Pi_l(p) = p$ for all $p \in P_l^n$ and hence the mapping $\Pi_l : C(E) \rightarrow P_l^n$ is a projection. Let

$$\|\Pi_l\| := \max_{x \in E} \sum_{i=1}^N |l_i(x)|.$$

A famous property says that for any $p \in P_l^n$,

$$\|p\|_\infty \leq \|\Pi_l\| \max_{1 \leq i \leq N} |p(x_i)|$$

The compact set E discussed in this paper will be mainly an n -dimensional simplex T_n whose definition can be found in [2].

It will be convenient to adopt **barycentric coordinates** when discussing points in a simplex. Suppose $v_1, \dots, v_{n+1}$ are the vertices of T_n . Then any $x \in T_n$ can be written as a convex combination of the vertices:

$$x = \sum_{i=1}^{n+1} c_i v_i$$

where $\sum_{i=1}^{n+1} c_i = 1$ and $c_i \geq 0$ for all i . The barycentric coordinate of x is then $(c_1, \dots, c_{n+1})$. Let's define "evenly spaced" points of **degree** l to be those points whose barycentric coordinates are of the form

$$(k_1/l, k_2/l, \dots, k_{n+1}/l), \quad k_i \text{ nonnegative integers with } \sum_{i=1}^{n+1} k_i = l.$$

Obviously the number of such points in T_n is exactly $N = \dim P_l^n$. Moreover, by [1], we know that evenly spaced points form a determining set for P_l^n .

The following lemma cited from [1] will be needed.

Lemma 1.1 *For the above evenly spaced points*

$$\|\Pi_l\| \leq \binom{2l-1}{l}.$$

Moreover, as $n \rightarrow \infty$, $\|\Pi_l\| \rightarrow \binom{2l-1}{l}$.

Now, we are going to prove a lemma which plays a crucial role in our construction of the error bound.

Lemma 1.2 *Let $Q \subseteq R^n$ be an n simplex in R^n and Y be the set of evenly spaced points of degree l in Q . Then, for any point x in Q , there is a measure σ supported on Y such that*

$$\int p(y) d\sigma(y) = p(x)$$

for all p in P_l^n , and

$$\int d|\sigma|(y) \leq \binom{2l-1}{l}.$$

Proof . Let $Y = \{y_1, \dots, y_N\}$ be the set of evenly spaced points of degree l in Q . Denote P_l^n by V . For any $x \in Q$, let δ_x be the point-evaluation functional. Define $T : V \rightarrow T(V) \subseteq R^N$ by $T(v) = (\delta_{y_i}(v))_{y_i \in Y}$. Then T is injective. Define $\tilde{\psi}$ on $T(V)$ by $\tilde{\psi}(w) = \delta_x(T^{-1}w)$. By the Hahn-Banach theorem, $\tilde{\psi}$ has a norm-preserving extension $\tilde{\psi}_{ext}$ to R^N . By the Riesz representation theorem, each linear functional on R^N can be represented by the inner product with a fixed vector. Thus, there exists $z \in R^N$ with

$$\tilde{\psi}_{ext}(w) = \sum_{j=1}^N z_j w_j$$

and $\|z\|_{(R^N)^*} = \|\tilde{\psi}_{ext}\|$. If we adopt the l_∞ -norm on R^N , the dual norm will be the l_1 -norm. Thus $\|z\|_{(R^N)^*} = \|z\|_1 = \|\tilde{\psi}_{ext}\| = \|\tilde{\psi}\| = \|\delta_x T^{-1}\|$.

Now, for any $p \in V$, by setting $w = T(p)$, we have

$$\delta_x(p) = \delta_x(T^{-1}w) = \tilde{\psi}(w) = \tilde{\psi}_{ext}(w) = \sum_{j=1}^N z_j w_j = \sum_{j=1}^N z_j \delta_{y_j}(p).$$

This gives

$$p(x) = \sum_{j=1}^N z_j p(y_j) \tag{5}$$

where $|z_1| + \dots + |z_N| = \|\delta_x T^{-1}\|$.

Note that

$$\|\delta_x T^{-1}\| = \sup_{\substack{w \in T(V) \\ w \neq 0}} \frac{\|\delta_x T^{-1}(w)\|}{\|w\|_{R^N}}$$

$$\begin{aligned}
 &= \sup_{\substack{w \in T(V) \\ w \neq 0}} \frac{|\delta_x p|}{\|T(p)\|_{R^N}} \\
 &\leq \sup_{\substack{p \in V \\ p \neq 0}} \frac{|p(x)|}{\max_{j=1,\dots,N} |p(y_j)|} \\
 &\leq \sup_{\substack{p \in V \\ p \neq 0}} \frac{\|\Pi_l\| \max_{j=1,\dots,N} |p(y_j)|}{\max_{j=1,\dots,N} |p(y_j)|} \\
 &= \|\Pi_l\| \\
 &\leq \binom{2l-1}{l}.
 \end{aligned}$$

Therefore $|z_1| + \dots + |z_N| \leq \binom{2l-1}{l}$ and our lemma follows immediately by (5). $\square$

1.2 Radial Functions and Borel Measures

Before moving on to our main result, some background for interpolation is necessary. First, the space of complex-valued functions on R^n that are compactly supported and infinitely differentiable is denoted by $\mathcal{D}$. The Fourier transform of a function ϕ in $\mathcal{D}$ is

$$\hat{\phi}(\xi) = \int e^{-i\langle x, \xi \rangle} \phi(x) dx.$$

Then a crucial lemma introduced in [5] but modified by Madych and Nelson in [9] says that for any continuous conditionally positive definite function h of order m , the Fourier transform of h uniquely determines a positive Borel measure μ on $R^n \setminus \{0\}$ and constants a_γ , $|\gamma| = 2m$ as follows: For all $\psi \in \mathcal{D}$

$$\begin{aligned}
 \int h(x) \psi(x) dx &= \int \left\{ \hat{\psi}(\xi) - \hat{\chi}(\xi) \sum_{|\gamma| < 2m} D^\gamma \hat{\psi}(0) \frac{\xi^\gamma}{\gamma!} \right\} d\mu(\xi) \\
 &\quad + \sum_{|\gamma| \leq 2m} D^\gamma \hat{\psi}(0) \frac{a_\gamma}{\gamma!},
 \end{aligned} \tag{6}$$

where for every choice of complex numbers c_α , $|\alpha| = m$,

$$\sum_{|\alpha|=m} \sum_{|\beta|=m} a_{\alpha+\beta} c_\alpha \bar{c}_\beta \geq 0.$$

Here χ is a function in $\mathcal{D}$ such that $1 - \hat{\chi}(\xi)$ has a zero of order $2m+1$ at $\xi = 0$; both of the integrals

$$\int_{0 < |\xi| < 1} |\xi|^{2m} d\mu(\xi), \quad \int_{|\xi| \geq 1} d\mu(\xi)$$

are finite. The choice of χ affects the value of the coefficients a_γ for $|\gamma| < 2m$.

2. MAIN RESULT

Before showing our main result, we need some lemmas. First, recall the famous formula of Stirling.

Stirling's Formula: $n! \sim \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$.

The approximation is very reliable even for small n . For example, when $n = 10$, the relative error is only 0.83% . The larger n is, the better the approximation is. For further details, we refer the reader to [3] and [4].

Lemma 2.1 Let $\rho_1 = \frac{1}{e}$ and $\rho_2 = \frac{3^{\frac{1}{6}}}{e} \sim \frac{1.2}{e}$. Then

$$\sqrt{2\pi\rho_1^k} k^k \leq k! \leq \sqrt{2\pi\rho_2^k} k^k$$

for all positive integer k .

Proof. Note that

$$\frac{1}{e}, \frac{\sqrt{2}}{e^2}, \frac{\sqrt{3}}{e^3}, \frac{\sqrt{4}}{e^4}, \frac{\sqrt{5}}{e^5}, \dots$$

can be expressed by

$$\frac{1}{e}, \left(\frac{2^{\frac{1}{4}}}{e}\right)^2, \left(\frac{3^{\frac{1}{6}}}{e}\right)^3, \left(\frac{4^{\frac{1}{8}}}{e}\right)^4, \left(\frac{5^{\frac{1}{10}}}{e}\right)^5, \dots$$

Now,

$$\sup \left\{ \frac{1}{e}, \frac{2^{\frac{1}{4}}}{e}, \frac{3^{\frac{1}{6}}}{e}, \frac{4^{\frac{1}{8}}}{e}, \frac{5^{\frac{1}{10}}}{e}, \dots \right\} = \frac{3^{\frac{1}{6}}}{e}$$

implies that $\frac{\sqrt{k}}{e^k} \leq \rho_2^k$ for all k . Thus $k! \sim \sqrt{2\pi} \frac{\sqrt{k}}{e^k} \cdot k^k \leq \sqrt{2\pi\rho_2^k} \cdot k^k$.

The remaining part $\sqrt{2\pi\rho_1^k} k^k \leq k!$ follows by observing that

$$\sqrt{2\pi} \left(\frac{1}{e}\right)^k k^k \leq \sqrt{2\pi} \left(\frac{1}{e}\right)^k \cdot \sqrt{k} \cdot k^k \sim k!.$$

□

Lemma 2.2 Let $\rho = \frac{\sqrt{3}}{e}$. Then $k! \leq \sqrt{2\pi\rho^k} k^{k-1}$ for all $k \geq 1$.

Proof. First,

$$k! \sim \sqrt{2\pi} \left(\frac{1}{e}\right)^k \cdot \sqrt{k} \cdot k^k = \sqrt{2\pi} \cdot \frac{k^{\frac{3}{2}}}{e^k} \cdot k^{k-1}.$$

Note that $\left\{ \frac{k^{\frac{3}{2}}}{e^k} : k = 1, 2, 3, \dots \right\}$ can be expressed by

$$\left\{ \frac{1}{e}, \frac{2^{\frac{3}{4}}}{e}, \frac{3^{\frac{1}{2}}}{e}, \frac{4^{\frac{3}{8}}}{e}, \dots \right\}.$$

Our lemma follows by noting that

$$\sup \left\{ \frac{1}{e}, \frac{2^{\frac{3}{4}}}{e}, \frac{3^{\frac{1}{2}}}{e}, \frac{4^{\frac{3}{8}}}{e}, \dots \right\} = \frac{\sqrt{3}}{e}.$$

□

Lemma 2.3 Let $h(x) = e^{-\beta|x|^2}$, $\beta > 0$, be the Gaussian function in R^n , and μ be the measure defined in (6). For any positive even integer l ,

$$\int_{R^n} |\xi|^l d\mu(\xi) \leq \pi^{\frac{n+1}{2}} \cdot n \cdot \alpha_n \cdot 2^{\frac{l+n+2}{2}} \cdot \rho^{\frac{l+n-1}{2}} \cdot \beta^{\frac{l}{2}} \cdot (l+n-1)^{\frac{l+n-3}{2}} \cdot \left(2 + \frac{1}{e}\right)$$

for odd n , and

$$\int_{R^n} |\xi|^l d\mu(\xi) \leq \pi^{\frac{n+1}{2}} \cdot n \cdot \alpha_n \cdot 2^{\frac{l+n+3}{2}} \cdot \rho^{\frac{l+n-2}{2}} \cdot \beta^{\frac{l}{2}} \cdot (l+n-2)^{\frac{l+n-4}{2}}$$

for even n , where $\rho = \frac{\sqrt{3}}{e}$ and α_n is the volume of the unit ball in R^n .

Proof.

$$\begin{aligned} & \int_{R^n} |\xi|^l d\mu(\xi) \\ &= \left(\frac{\pi}{\beta}\right)^{\frac{n}{2}} \int_{R^n} \frac{|\xi|^l}{e^{\frac{|\xi|^2}{4\beta}}} d\xi \\ &= \left(\frac{\pi}{\beta}\right)^{\frac{n}{2}} \cdot n \cdot \alpha_n \cdot \int_0^\infty \frac{r^l \cdot r^{n-1}}{e^{\frac{r^2}{4\beta}}} dr \\ &= \left(\frac{\pi}{\beta}\right)^{\frac{n}{2}} \cdot n \cdot \alpha_n \cdot (2\sqrt{\beta})^{l+n} \cdot \int_0^\infty \frac{r^{l+n-1}}{e^{r^2}} dr \\ &= \begin{cases} \pi^{\frac{n}{2}} \cdot n \cdot \alpha_n \cdot 2^{n-1+l} \cdot \beta^{\frac{l}{2}} \cdot \left(\frac{l+n-2}{2}\right) \left(\frac{l+n-2}{2} - 1\right) \cdots \left(\frac{1}{2}\right) \int_0^\infty \frac{1}{\sqrt{ve^v}} dv \\ \text{if } n \text{ is odd,} \\ \pi^{\frac{n}{2}} \cdot n \cdot \alpha_n \cdot 2^{n-1+l} \cdot \beta^{\frac{l}{2}} \left(\frac{l+n-2}{2}\right)! \text{ if } n \text{ is even.} \end{cases} \end{aligned}$$

Let $b = \lceil \frac{l+n+2}{2} \rceil$. Then $b! \leq \sqrt{2\pi} \rho^b \cdot b^{b-1}$. Then for odd n , $b = \frac{l+n+1}{2}$. Thus

$$b! \leq \sqrt{2\pi} \rho^{\frac{l+n-1}{2}} \cdot \left(\frac{l+n-1}{2}\right)^{\frac{l+n-3}{2}} = \sqrt{2\pi} \cdot 2^{\frac{-l-n+3}{2}} \cdot \rho^{\frac{l+n-1}{2}} \cdot (l+n-1)^{\frac{l+n-3}{2}},$$

and

$$\int_{R^n} |\xi|^l d\mu(\xi) \leq \pi^{\frac{n+1}{2}} \cdot n \cdot \alpha_n \cdot \left(2 + \frac{1}{e}\right) \cdot 2^{\frac{l+n+2}{2}} \cdot \beta^{\frac{l}{2}} \cdot \rho^{\frac{l+n-1}{2}} \cdot (l+n-1)^{\frac{l+n-3}{2}},$$

by noting that $\int_0^\infty \frac{1}{\sqrt{ve^v}} dv \leq 2 + \frac{1}{e}$. For even n , $b = \frac{l+n-2}{2}$. Thus

$$b! \leq \sqrt{2\pi} \rho^{\frac{l+n-2}{2}} \cdot \left(\frac{l+n-2}{2}\right)^{\frac{l+n-4}{2}} = \sqrt{2\pi} \cdot 2^{\frac{-l-n+4}{2}} \cdot \rho^{\frac{l+n-2}{2}} \cdot (l+n-2)^{\frac{l+n-4}{2}},$$

and

$$\int_{R^n} |\xi|^l d\mu(\xi) \leq \pi^{\frac{n+1}{2}} \cdot n \cdot \alpha_n \cdot 2^{\frac{l+n+3}{2}} \cdot \beta^{\frac{l}{2}} \cdot \rho^{\frac{l+n-2}{2}} \cdot (l+n-2)^{\frac{l+n-4}{2}}. \quad \square$$

Our interpolation is based on a function space called **native space**, denoted by $C_{h,m}$. If

$$\mathcal{D}_m = \left\{ \phi \in \mathcal{D} : \int x^\alpha \phi(x) dx = 0 \text{ for all } |\alpha| < m \right\}$$

, then $C_{h,m}$ is the class of those continuous functions f which satisfy

$$\left| \int f(x) \phi(x) dx \right| \leq c(f) \left\{ \int h(x-y) \phi(x) \overline{\phi(y)} dx dy \right\}^{\frac{1}{2}} \quad (7)$$

for some constant $c(f)$ and all ϕ in $\mathcal{D}_m$. If $f \in C_{h,m}$, let $\|f\|_h$ denotes the smallest constant $c(f)$ for which (7) is true. It can be shown that $\|f\|_h$ is a semi-norm and $C_{h,m}$ is a semi-Hilbert space; in the case $m = 0$ it is a norm and a Hilbert space respectively. In this paper both the interpolating and interpolated functions belong to the native space.

The function space $C_{h,m}$ was introduced by Madych and Nelson in [8] and [9]. Later Luh made a lucid characterization in [6] and [7]. Although there is an equivalent expression for $C_{h,m}$ which is easier to understand, we still adopt Madych and Nelson's definition to show the author's respect for them. The main result of this paper is the following theorem.

Theorem 2.4 Let $h(x) := e^{-\beta|x|^2}$ be the Gaussian function in R^n . For any positive number b_0 , there are positive constants δ_0, c_1, c_2 , and c_3 independent of n , for which the following is true: If $f \in C_{h,m}$, the native space induced by h , and s is the h spline that interpolates f on a subset X of R^n , then

$$|f(x) - s(x)| \leq c_1 \sqrt{\delta} (c_2 \delta)^{\frac{c_3}{\delta}} \cdot \|f\|_h \quad (8)$$

for all x in a subset Ω of R^n , and $0 < \delta \leq \delta_0$, where Ω satisfies the property that for any x in Ω and any number $\frac{b_0}{2} \leq r \leq b_0$, there is an n simplex Q with diameter $\text{diam} Q = r$, $x \in Q \subseteq \Omega$, such that for any integer l with $\frac{b_0}{\delta} \leq l \leq \frac{2b_0}{\delta}$, there is on Q an evenly spaced set of centers from X of degree $l-1$. (In fact, the set X can be chosen to consist of these evenly spaced centers in Q only.) Here $\|f\|_h$ is the h -norm of f in the native space. The numbers δ_0, c_1, c_2 , and c_3 are given by $\delta_0 := \min \left\{ b_0, \frac{1}{\rho_3^4 \cdot 3^3 \cdot 2^7 \cdot b_0^3} \right\}$ where $\rho_3 = 12^{\frac{1}{4}} \cdot \sqrt{e\beta}$;

$$\begin{cases} c_1 := \begin{cases} \Delta'' \cdot \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{b_0}} & \text{for odd } n, \\ \Delta'' \cdot \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{b_0}} & \text{for even } n, \end{cases} \\ c_2 := \rho_3^4 \cdot 3^3 \cdot 2^7 \cdot b_0^3, \\ c_3 := \frac{b_0}{4}, \end{cases}$$

where Δ'' is defined by

$$\Delta'' := \begin{cases} \sqrt{2 + \frac{1}{e}} \cdot \pi^{\frac{n-1}{4}} \cdot (n\alpha_n)^{\frac{1}{2}} \cdot 2^{\frac{n}{4}} \cdot \rho^{\frac{n-1}{4}} & \text{for odd } n, \\ \pi^{\frac{n-1}{4}} \cdot (n\alpha_n)^{\frac{1}{2}} \cdot 2^{\frac{n+1}{4}} \cdot \rho^{\frac{n-2}{4}} & \text{for even } n, \end{cases} \text{ with } \rho = \frac{\sqrt{3}}{e} \text{ as in Lemma 2.3}$$

, where the number α_n denotes the volume of the unit ball in R^n .

In particular, if the point x in Ω is fixed, the only requirement for Ω is the existence of an n simplex Q , with $\text{diam} Q = r$, $x \in Q \subseteq \Omega$, satisfying the afore-mentioned property of evenly spaced centers.

Proof. For any $b_0 > 0$, let $\delta_0 := \min \left\{ b_0, \frac{1}{\rho_3^4 \cdot 3^3 \cdot 2^7 \cdot b_0^3} \right\}$ where $\rho_3 = 12^{\frac{1}{4}} \cdot \sqrt{e\beta}$. For any $0 < \delta \leq \delta_0$, there exists an integer l such that $1 \leq \frac{\delta}{b_0} l \leq 2$ since $0 < \frac{\delta}{b_0} \leq 1$. Such l satisfies $\frac{b_0}{\delta} \leq l \leq \frac{2b_0}{\delta}$ and $\frac{b_0}{2} \leq \frac{\delta}{2} l \leq b_0$.

For any $x \in \Omega$, let Q be an n simplex containing x such that $Q \subseteq \Omega$ and has diameter $\text{diam} Q = \frac{\delta l}{2}$. Then Theorem 4.2 of [9] implies that

$$|f(x) - s(x)| \leq c_l \|f\|_h \int_{R^n} |y - x|^l d|\sigma|(y) \quad (9)$$

whenever $l > 0$, where σ is any measure supported on X such that

$$\int_{R^n} p(y) d\sigma(y) = p(x) \quad (10)$$

for all polynomials p in P_{l-1}^n . Here

$$c_l = \left\{ \int_{R^n} \frac{|\xi|^{2l}}{(l!)^2} d\mu(\xi) \right\}^{\frac{1}{2}}$$

whenever $l > 0$. Be careful. We temporarily use c_1, c_2, c_3 to denote numbers which are totally different from the c_1, c_2, c_3 mentioned in the theorem.

Let Y be the set of evenly spaced centers from X of degree $l-1$ on Q . By Lemma1.2, there is a measure σ supported on Y such that (10) is satisfied and

$$\int d|\sigma|(y) \leq \binom{2l-3}{l-1}$$

The crux of our error estimate is to bound

$$I = c_l \int_{R^n} |y-x|^l d|\sigma|(y).$$

Now Lemma2.3 applies. For odd n ,

$$\begin{aligned} c_l &= \frac{1}{l!} \left\{ \int_{R^n} |\xi|^{2l} d\mu(\xi) \right\}^{1/2} \\ &\leq \frac{1}{l!} \left\{ \pi^{\frac{n+1}{2}} \cdot n \cdot \alpha_n \cdot 2^{\frac{2l+n-2}{2}} \cdot \rho^{\frac{2l+n-1}{2}} \cdot \beta^l \cdot (2l+n-1)^{\frac{2l+n-3}{2}} \cdot \left(2 + \frac{1}{e}\right) \right\}^{1/2} \\ &\leq \frac{1}{l!} \left\{ \pi^{\frac{n+1}{4}} \cdot (n\alpha_n)^{1/2} \cdot 2^{\frac{n+2}{4}} \cdot \rho^{\frac{n-1}{4}} \cdot (\sqrt{2\rho\beta})^l \cdot (2l+n-1)^{\frac{2l+n-3}{4}} \cdot \sqrt{2 + \frac{1}{e}} \right\} \\ &\leq \left\{ \pi^{\frac{n+1}{4}} (n\alpha_n)^{1/2} \cdot 2^{\frac{n+2}{4}} \cdot \rho^{\frac{n-1}{4}} \cdot (\sqrt{2\rho\beta})^l (2l+n-1)^{\frac{2l+n-3}{4}} \left(2 + \frac{1}{e}\right) \right\} / \left\{ \sqrt{2\pi} \cdot \rho_1^l \cdot l^l \right\} \\ &\quad \text{by Lemma2.1} \\ &= \pi^{\frac{n-1}{4}} \cdot (n\alpha_n)^{1/2} \cdot 2^{\frac{n}{4}} \cdot \rho^{\frac{n-1}{4}} \cdot \rho_3^l \cdot \left(\frac{1}{l!}\right) (2l+n-1)^{\frac{2l+n-3}{4}} \cdot \left(2 + \frac{1}{e}\right)^{1/2} \\ &\quad \text{where } \rho_3 = \frac{\sqrt{2\rho\beta}}{\rho_1} \\ &= \Delta'' \cdot \rho_3^l \cdot \frac{1}{l!} (2l+n-1)^{\frac{2l+n-3}{4}} \text{ where } \Delta'' = \sqrt{2 + \frac{1}{e}} \cdot \pi^{\frac{n-1}{4}} (n\alpha_n)^{1/2} 2^{n/4} \rho^{\frac{n-1}{4}}. \end{aligned}$$

Note that the diameter of Q is $\frac{\delta l}{2}$. This gives for $l \geq n-3$

$$\begin{aligned} I &\leq \Delta'' \rho_3^l l^{-l} (2l+n-1)^{\frac{2l+n-3}{4}} (\text{diam} Q)^l \binom{2l-3}{l-1} \\ &\leq \Delta'' \rho_3^l l^{-l} (3l)^{\frac{3l}{4}} \left(\frac{\delta}{2} l\right)^l \binom{2l-3}{l-1} \text{ if } l \geq n-3 \\ &= \Delta'' \rho_3^l (3^{3/4})^l \left(\frac{\delta}{2} l^{3/4}\right)^l \binom{2l-3}{l-1} \text{ if } l \geq n-3 \\ &= \Delta'' (\rho_3 3^{3/4} \frac{\delta}{2} l^{3/4})^l \binom{2l-3}{l-1} \\ &\leq \Delta'' \left\{ \rho_3 3^{3/4} \cdot \frac{1}{2} \cdot \delta \left(\frac{2b_0}{\delta}\right)^{3/4} \right\}^l \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{l-1}} 4^l \cdot \frac{1}{4} \text{ by Stirling's formula and} \\ &\quad \frac{b_0}{\delta} \leq l \leq \frac{2b_0}{\delta} \\ &= \Delta'' \left\{ \rho_3 3^{3/4} \cdot 2(2b_0)^{3/4} \delta^{1/4} \right\}^l \frac{1}{\sqrt{16\pi}} \frac{1}{\sqrt{l-1}} \\ &\leq \Delta'' \left\{ \rho_3 3^{3/4} \cdot 2 \cdot 2^{3/4} \cdot b_0^{3/4} \cdot \delta^{1/4} \right\}^{\frac{b_0}{\delta}} \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l-1}} \end{aligned}$$

$$\begin{aligned}
 & \text{where } \rho_3 \cdot 3^{3/4} \cdot 2 \cdot 2^{3/4} \cdot b_0^{3/4} \cdot \delta^{1/4} \leq 1 \text{ iff } \delta^{1/4} \leq \frac{1}{\rho_3 3^{3/4} \cdot 2^{7/4} \cdot b_0^{3/4}}, \text{ which} \\
 & \text{is guaranteed by } \delta \leq \delta_0 := \min \left\{ b_0, \frac{1}{\rho_3^4 \cdot 3^3 \cdot 2^7 \cdot b_0^3} \right\} \\
 & \leq \Delta'' \left\{ \rho_3^4 \cdot 3^3 \cdot 2^7 \cdot b_0^3 \cdot \delta \right\}^{\frac{b_0}{46}} \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l-1}} \\
 & \leq \Delta'' \cdot \frac{1}{\sqrt{16\pi}} \cdot \{G\delta\}^{\frac{g}{\delta}} \cdot \frac{1}{\sqrt{l-1}} \text{ where } G = \rho_3^4 3^3 2^7 b_0^3 \text{ and } g = \frac{b_0}{4} \\
 & \sim \frac{\Delta''}{\sqrt{16\pi}} \{G\delta\}^{\frac{g}{\delta}} \frac{1}{\sqrt{l}} \text{ (if } l \gg 0) \\
 & \leq \frac{\Delta''}{\sqrt{16\pi}} \sqrt{\frac{1}{b_0}} \sqrt{\delta} \{G\delta\}^{\frac{g}{\delta}} \text{ since } \frac{b_0}{\delta} \leq l \leq \frac{2b_0}{\delta} \text{ if } l \geq n-3
 \end{aligned}$$

For even n , if $l \geq n-4$,

$$\begin{aligned}
 I &= c_l \int_{R^n} |y-x|^l d|\sigma|(y) \\
 &\leq \Delta'' \rho_3^l \cdot l^{-l} (2l+n-2)^{\frac{2l+n-4}{4}} (\dim Q)^l \binom{2l-3}{l-1} \\
 &\quad \text{where } \Delta'' = \pi^{\frac{n-1}{4}} \cdot (n\alpha_n)^{\frac{1}{2}} \cdot 2^{\frac{n+1}{4}} \cdot \rho^{\frac{n-2}{4}} \text{ and } \rho_3 = \frac{\sqrt{2\rho\beta}}{\rho_1} \\
 &= \Delta'' \rho_3^l \cdot l^{-l} (2l+n-2)^{\frac{2l+n-4}{4}} \left(\frac{\delta l}{2} \right)^l \binom{2l-3}{l-1} \\
 &\leq \Delta'' \rho_3^l \cdot (3l)^{\frac{3l}{4}} \cdot \delta^l \cdot \left(\frac{1}{2} \right)^l \binom{2l-3}{l-1} \text{ since } l \geq n-4 \\
 &\leq \Delta'' \left(\rho_3 3^{\frac{3}{4}} l^{\frac{3}{4}} \frac{1}{2} \delta \right)^l \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{l-1}} 4^{l-1} \text{ by Stirling's formula} \\
 &= \Delta'' \frac{1}{\sqrt{\pi}} \cdot \frac{1}{4} \left(\rho_3 3^{\frac{3}{4}} \cdot \frac{1}{2} \cdot \delta l^{\frac{3}{4}} 4 \right)^l \frac{1}{\sqrt{l-1}} \\
 &\sim \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l}} (\rho_3 \cdot 2 \cdot 3^{\frac{3}{4}} \delta l^{\frac{3}{4}})^l \text{ (if } l \gg 0) \\
 &\leq \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l}} \left\{ \rho_3 \cdot 2 \cdot 3^{\frac{3}{4}} \cdot \delta \cdot \left(\frac{2b_0}{\delta} \right)^{\frac{3}{4}} \right\}^l \text{ since } \frac{b_0}{\delta} \leq l \leq \frac{2b_0}{\delta} \\
 &= \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l}} \left\{ \rho_3 \cdot 2^{\frac{7}{4}} \cdot 3^{\frac{3}{4}} \cdot b_0^{\frac{3}{4}} \cdot \delta^{\frac{1}{4}} \right\}^l \\
 &\leq \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l}} \left\{ \rho_3 \cdot 2^{\frac{7}{4}} \cdot 3^{\frac{3}{4}} \cdot b_0^{\frac{3}{4}} \cdot \delta^{\frac{1}{4}} \right\}^{\frac{b_0}{\delta}} \text{ where} \\
 &\quad \rho_3 \cdot 2^{\frac{7}{4}} \cdot 3^{\frac{3}{4}} \cdot b_0^{\frac{3}{4}} \cdot \delta^{\frac{1}{4}} \leq 1 \text{ iff } \delta \leq \frac{1}{\rho_3^4 \cdot 2^7 \cdot 3^3 \cdot b_0^3}, \text{ which is guaranteed by} \\
 &\quad \delta \leq \delta_0 := \min \left\{ \frac{1}{\rho_3^4 \cdot 2^7 \cdot 3^3 \cdot b_0^3}, b_0 \right\} \\
 &\leq \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l}} \left\{ \rho_3^4 \cdot 2^7 \cdot 3^3 \cdot b_0^3 \cdot \delta \right\}^{\frac{b_0}{46}}
 \end{aligned}$$

$$\begin{aligned}
 &= \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{l}} \{G\delta\}^{\frac{g}{\delta}} \text{ where } G = \rho_3^4 \cdot 2^7 \cdot 3^3 \cdot b_0^3 \text{ and } g = \frac{b_0}{4} \\
 &\leq \Delta'' \frac{1}{\sqrt{16\pi}} \cdot \frac{1}{\sqrt{b_0}} \cdot \sqrt{\delta} \{G\delta\}^{\frac{g}{\delta}} \text{ since } \frac{b_0}{\delta} \leq l \leq \frac{2b_0}{\delta}.
 \end{aligned}$$

Our error bound (8) then follows immediately by letting $\rho_3 = \sqrt{e\beta} \cdot 12^{\frac{1}{4}}$. $\square$

In the proof of Theorem 2.4 although we require that $l \geq n-3$ for n odd and $l \geq n-4$ for n even, besides $l \gg 0$, it usually causes no trouble because $\delta \rightarrow 0$ implies $l \rightarrow \infty$ and usually δ is very small. In order to make the theorem easier to understand, we don't put these requirements into the theorem.

Remark. In theorem 2.4 we avoid using the well-known term “fill distance” for scattered data approximation because in our approach the data points are not purely scattered. However the number δ is in spirit equivalent to the fill distance in the sense that $\delta \rightarrow 0$ iff the fill distance $d(Q, Y) \rightarrow 0$ where Y is the evenly spaced centers from X of degree $l-1$ on Q . Note that $d(Q, Y) \rightarrow 0$ iff $l \rightarrow \infty$, and $l \rightarrow \infty$ iff $\delta \rightarrow 0$.

Furthermore, although the function space $C_{h,m}$ is defined in a complicated way, there is a simple expression for it. This can be seen in Wendland's book [11] where Madych and Nelson's native space [8] and Wu and Schaback's native space [12] are unified in an elegant way.

3. COMPARISON

The exponential-type error bound for Gaussian interpolation raised by Madych and Nelson in [10] is of the form

$$|f(x) - s(x)| \leq a_1(a_2\delta)^{\frac{a_3}{\delta}} \|f\|_h, \text{ as } \delta \rightarrow 0 \quad (11)$$

where a_1 is about the same as c_1 of (8), $a_3 = \frac{b_0}{8\gamma_n}$ and $a_2 = (3^{3/4} \cdot e \cdot \sqrt{2\rho\beta} \cdot \sqrt{n} \cdot e^{2n\gamma_n})^4 \cdot b_0^3 \cdot \gamma_n$ where $\rho = \frac{\sqrt{3}}{e}$, b_0 is the side length of a cube and γ_n is defined recursively by

$$\gamma_1 = 2, \gamma_n = 2n(1 + \gamma_{n-1}) \text{ if } n > 1.$$

These can be seen in [10] and [?]. Note that $\gamma_n \rightarrow \infty$ rapidly as $n \rightarrow \infty$. The first few examples are

$$\gamma_1 = 2, \gamma_2 = 12, \gamma_3 = 78, \gamma_4 = 632 \text{ and } \gamma_5 = 6330.$$

This means that for high dimensions, a_2 will become extremely large and a_3 extremely small, making the error bound meaningless.

Our new approach avoids this drawback. Although we require that the centers be evenly spaced in the simplex, it causes no trouble at all both theoretically and practically.

REFERENCES

- [1] L.P. Bos. (1983). Bounding the Lebesgue Function for Lagrange Interpolation in a Simplex. *J. of Approx. Theory*, 38, 43-59.
- [2] W. Fleming. (1977). *Functions of Several Variables*, Second Edition. Springer-Verlag.
- [3] J. von zur Gathen and J. Gerhard. (2003). *Modern Computer Algebra*, Second Edition. Cambridge University Press.
- [4] R.L. Graham, D.E. Knuth, and O. Patashnik. (1994). *Concrete Mathematics*. Addison-Wesley, Reading MA, 2nd edition.

- [5] I.M. Gelfand and N.Y. Vilenkin. (1964). *Generalized Functions*, 4. New York: Academics Press.
- [6] L.T. Luh. (2001). The Equivalence Theory of Native Spaces. *Approx. Theory Appl.*, 17(1), 76-96.
- [7] L.T. Luh,(2008). The Crucial Constants in the Exponential-type Error Estimates for Gaussian Interpolation. *Analysis in Theory Appl.*, 24(2), 183–194.
- [8] W.R. Madych and S.A. Nelson. (1988). Multivariate interpolation and conditionally positive definite function. *Approx. Theory Appl.* 4(4), 77-89.
- [9] W.R. Madych and S.A. Nelson. (1990). Multivariate interpolation and conditionally positive definite function, II. *Math. Comp.* 54, 211-230.
- [10] W.R. Madych and S.A. Nelson. (1992). Bounds on Multivariate Polynomials and Exponential Error Estimates for Multivariate Interpolation. *J. Approx. Theory*, 70, 94-114.
- [11] H. Wendland. (2005). *Scattered Data Approximation*. Cambridge University Press.
- [12] Z. Wu and R. Schaback. (1993). Local Error Estimates for Radial Basis Function Interpolation of Scattered Data. *IMA J. of Numerical Analysis*.